



BIKTARVY®

bictegravir 50mg/emtricitabine 200mg/
tenofovir alafenamide 25mg tablets

(bik-TAR-vee)

MOST IMPORTANT INFORMATION ABOUT BIKTARVY

BIKTARVY may cause serious side effects, including:

- **Worsening of hepatitis B (HBV) infection.** If you have both HIV-1 and HBV, your HBV may suddenly get worse if you stop taking BIKTARVY. Do not stop taking BIKTARVY without first talking to your healthcare provider, as they will need to check your health regularly for several months.

ABOUT BIKTARVY

BIKTARVY is a complete, 1-pill, once-a-day prescription medicine used to treat HIV-1 in adults. It can either be used in people who have never taken HIV-1 medicines before, or people who are replacing their current HIV-1 medicines and whose healthcare provider determines they meet certain requirements.

BIKTARVY does not cure HIV-1 or AIDS. HIV-1 is the virus that causes AIDS.

Do NOT take BIKTARVY if you also take a medicine that contains:

- dofetilide
- rifampin
- any other medicines to treat HIV-1

BEFORE TAKING BIKTARVY

Tell your healthcare provider all your medical conditions, including if you:

- Have or have had any kidney or liver problems, including hepatitis infection.
- Are pregnant or plan to become pregnant.
- Are breastfeeding (nursing) or plan to breastfeed. Do not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, antacids, laxatives, vitamins, and herbal supplements, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that interact with BIKTARVY.

IMPORTANT FACTS

This is only a brief summary of important information about BIKTARVY® and does not replace talking to your healthcare provider about your condition and your treatment.

POSSIBLE SIDE EFFECTS OF BIKTARVY

BIKTARVY can cause serious side effects, including:

- Those in the “Most Important Information About BIKTARVY” section.
- Changes in your immune system.
- New or worse kidney problems, including kidney failure.
- Too much lactic acid in your blood (lactic acidosis), which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.
- Severe liver problems, which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark “tea-colored” urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.
- **The most common side effects of BIKTARVY** in clinical studies were diarrhea (6%), nausea (6%), and headache (5%).

These are not all the possible side effects of BIKTARVY. Tell your healthcare provider right away if you have any new symptoms while taking BIKTARVY.

Your healthcare provider will need to do tests to monitor your health before and during treatment with BIKTARVY.

HOW TO TAKE BIKTARVY

Take BIKTARVY 1 time each day with or without food.

GET MORE INFORMATION

- This is only a brief summary of important information about BIKTARVY. Talk to your healthcare provider or pharmacist to learn more.
- Go to BIKTARVY.com or call 1-800-GILEAD-5
- If you need help paying for your medicine, visit BIKTARVY.com for program information.



BIKTARVY, the BIKTARVY Logo, GILEAD, and the GILEAD Logo are trademarks of Gilead Sciences, Inc., or its related companies. Version date: December 2018

© 2019 Gilead Sciences, Inc. All rights reserved. BVYC0113 01/19