

(des-KOH-vee)

IMPORTANT FACTS

This is only a brief summary of important information about **DESCOVY**[®] and does not replace talking to your healthcare provider about your condition and your treatment.

MOST IMPORTANT INFORMATION ABOUT DESCOVY

DESCOVY may cause serious side effects, including:

- **Worsening of hepatitis B (HBV) infection.** DESCOVY is not approved to treat HBV. If you have both HIV-1 and HBV, your HBV may suddenly get worse if you stop taking DESCOVY. Do not stop taking DESCOVY without first talking to your healthcare provider, as they will need to check your health regularly for several months.

ABOUT DESCOVY

- DESCOVY is a prescription medicine that is used together with other HIV-1 medicines to treat HIV-1 in people who weigh at least 77 lbs (35kg). DESCOVY is **not** for use to help reduce the risk of getting HIV-1 infection.
- **DESCOVY does not cure HIV-1 or AIDS.** Ask your healthcare provider about how to prevent passing HIV-1 to others.

BEFORE TAKING DESCOVY

Tell your healthcare provider if you:

- Have or had any kidney or liver problems, including hepatitis infection.
- Have any other medical condition.
- Are pregnant or plan to become pregnant.
- Are breastfeeding (nursing) or plan to breastfeed. Do not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that should not be taken with DESCOVY.

HOW TO TAKE DESCOVY

- DESCOVY is a one pill, once a day HIV-1 medicine that is taken with other HIV-1 medicines.
- Take DESCOVY with or without food.

POSSIBLE SIDE EFFECTS OF DESCOVY

DESCOVY can cause serious side effects, including:

- Those in the “Most Important Information About DESCOVY” section.
- Changes in your immune system.
- New or worse kidney problems, including kidney failure.
- Too much lactic acid in your blood (lactic acidosis), which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.
- Severe liver problems, which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark “tea-colored” urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.

The most common side effect of DESCOVY is nausea.

These are not all the possible side effects of DESCOVY. Tell your healthcare provider right away if you have any new symptoms while taking DESCOVY.

Your healthcare provider will need to do tests to monitor your health before and during treatment with DESCOVY.

GET MORE INFORMATION

- This is only a brief summary of important information about DESCOVY. Talk to your healthcare provider or pharmacist to learn more.
- Go to DESCOVY.com or call 1-800-GILEAD-5
- If you need help paying for your medicine, visit DESCOVY.com for program information.