

(des-KOH-vee)

## MOST IMPORTANT INFORMATION ABOUT DESCOVY FOR PrEP

### Before starting and while taking DESCOVY for PrEP:

- **You must be HIV-1 negative.** You must get tested to make sure that you do not already have HIV-1. Do not take DESCOVY for PrEP to reduce the risk of getting HIV-1 unless you are confirmed to be HIV-1 negative. Get tested for HIV-1 immediately before and at least every 3 months while taking DESCOVY.
- **Many HIV-1 tests can miss HIV-1 infection in a person who has recently become infected.** Symptoms of new HIV-1 infection include tiredness, fever, joint or muscle aches, headache, sore throat, vomiting, diarrhea, rash, night sweats, and/or enlarged lymph nodes in the neck or groin. Tell your healthcare provider if you have had a flu-like illness within the last month before starting or while taking DESCOVY.
- **DESCOVY does not prevent other sexually transmitted infections (STIs).** Practice safer sex by using a latex or polyurethane condom to reduce the risk of getting STIs.
- **DESCOVY by itself is not a complete treatment for HIV-1.** If you do become HIV-1 positive, you need more medicine than DESCOVY alone to treat HIV-1.

### DESCOVY may cause serious side effects, including:

- **Worsening of hepatitis B (HBV) infection.** Your healthcare provider will test you for HBV. If you have HBV, your HBV may suddenly get worse if you stop taking DESCOVY. Do not stop taking DESCOVY without first talking to your healthcare provider, as they will need to check or give you HBV medicine.

## ABOUT DESCOVY FOR PrEP

DESCOVY for PrEP is a once-daily prescription medicine to reduce the risk of getting HIV-1 infection in adults and adolescents who weigh at least 77 pounds. It is not known if DESCOVY is effective in reducing the risk of getting HIV-1 from certain types of sex.

- DESCOVY for PrEP is not for use in people assigned female at birth who are at risk of getting HIV-1 infection from vaginal sex, because its effectiveness has not been studied.
- To help determine your risk of getting HIV-1, talk openly with your healthcare provider about your sexual health.

### Do NOT take DESCOVY for PrEP if you:

- **Already have HIV-1 infection or if you do not know your HIV-1 status.**

## BEFORE TAKING DESCOVY FOR PrEP

### Tell your healthcare provider if you:

- Have or have had any kidney or liver problems, including hepatitis.
- Have any other medical conditions.

### Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that should not be taken with DESCOVY for PrEP.

## IMPORTANT FACTS

This is only a brief summary of important information about taking DESCOVY for PrEP™ (pre-exposure prophylaxis). This does not replace talking to your healthcare provider about your medicine.

## POSSIBLE SIDE EFFECTS OF DESCOVY FOR PrEP

### DESCOVY<sup>®</sup> can cause serious side effects, including:

- Those in the “Most Important Information About DESCOVY for PrEP” section.
- New or worse kidney problems, including kidney failure.
- Too much lactic acid in your blood (lactic acidosis), which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.
- Severe liver problems, which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark “tea-colored” urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.

**Common side effects** in people taking DESCOVY for PrEP include diarrhea, nausea, headache, fatigue, and stomach pain.

These are not all the possible side effects of DESCOVY. Tell your healthcare provider right away if you have any new symptoms while taking DESCOVY for PrEP.

**Your healthcare provider will need to do tests to monitor your health before and during treatment with DESCOVY for PrEP.**

## HOW TO TAKE DESCOVY FOR PrEP

- **Take 1 tablet once a day, every day**, not just when you think you have been exposed to HIV-1.
- Do not miss any doses. Missing doses may increase your risk of getting HIV-1 infection.
- Get tested for HIV-1 at least every 3 months. You must stay HIV-negative to keep taking DESCOVY for PrEP.

## HOW TO FURTHER REDUCE YOUR RISK

- Know your HIV status and the HIV status of your partners.
- Get tested for other STIs. Some STIs make it easier for HIV-1 to infect you.
- Practice safer sex by using latex or polyurethane condoms.
- Talk to your healthcare provider about all the ways to help reduce HIV risk.

## GET MORE INFORMATION

- This is only a brief summary of important information about DESCOVY for PrEP. Talk to your healthcare provider or pharmacist to learn more.
- Go to [DESCOVY.com/prep](https://www.descovy.com/prep) or call 1-800-GILEAD-5
- If you need help paying for your medicine, visit [DESCOVY.com/prep](https://www.descovy.com/prep) for program information.