

(jen-VOY-uh)

MOST IMPORTANT INFORMATION ABOUT GENVOYA

GENVOYA may cause serious side effects, including:

- **Worsening of hepatitis B (HBV) infection.** If you have both HIV-1 and HBV, your HBV may suddenly get worse if you stop taking GENVOYA. Do not stop taking GENVOYA without first talking to your healthcare provider, as they will need to check your health regularly for several months.

ABOUT GENVOYA

- GENVOYA is a prescription medicine used to treat HIV-1 in people 12 years of age and older who weigh at least 77 pounds and have never taken HIV-1 medicines before. GENVOYA can also be used to replace current HIV-1 medicines for some people who have an undetectable viral load (less than 50 copies/mL of virus in their blood), and have been on the same HIV-1 medicines for at least 6 months and have never failed HIV-1 treatment, and whose healthcare provider determines that they meet certain other requirements.
- **GENVOYA does not cure HIV-1 or AIDS.** HIV-1 is the virus that causes AIDS.

Do NOT take GENVOYA if you:

- Take a medicine that contains: alfuzosin (Uroxatral[®]), carbamazepine (Carbatrol[®], Carnexiv[®], Eptol[®], Equetro[®], Tegretol[®], Tegretol-XR[®], Teril[®]), cisapride (Propulsid[®], Propulsid Quicksolv[®]), dihydroergotamine (D.H.E. 45[®], Migranal[®]), ergotamine (Cafergot[®], Migergot[®], Ergostat[®], Medihaler Ergotamine[®], Wigraine[®], Wigrettes[®]), lomitapide (Juxtapid[®]), lovastatin (Altoprev[®], Mevacor[®]), lurasidone (Latuda[®]), midazolam (when taken by mouth), phenobarbital (Luminal[®]), phenytoin (Dilantin[®], Dilantin-125[®], Phenytek[®]), pimozone (Orap[®]), rifampin (Rifadin[®], Rifamate[®], Rifater[®], Rimactane[®]), sildenafil when used for lung problems (Revatio[®]), simvastatin (Vytorin[®], Zocor[®]), or triazolam (Halcion[®]).
- Take the herbal supplement St. John's wort.
- Take any other HIV-1 medicines at the same time.

GET MORE INFORMATION

- This is only a brief summary of important information about GENVOYA. Talk to your healthcare provider or pharmacist to learn more.
- Go to GENVOYA.com or call 1-800-GILEAD-5
- If you need help paying for your medicine, visit GENVOYA.com for program information.

IMPORTANT FACTS

This is only a brief summary of important information about GENVOYA[®] and does not replace talking to your healthcare provider about your condition and your treatment.

POSSIBLE SIDE EFFECTS OF GENVOYA

GENVOYA can cause serious side effects, including:

- Those in the "Most Important Information About GENVOYA" section.
- Changes in your immune system.
- New or worse kidney problems, including kidney failure.
- Too much lactic acid in your blood (lactic acidosis), which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.
- Severe liver problems, which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark "tea-colored" urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.

The most common side effect of GENVOYA is nausea.

These are not all the possible side effects of GENVOYA. Tell your healthcare provider right away if you have any new symptoms while taking GENVOYA.

Your healthcare provider will need to do tests to monitor your health before and during treatment with GENVOYA.

BEFORE TAKING GENVOYA

Tell your healthcare provider if you:

- Have or have had any kidney or liver problems, including hepatitis infection.
- Have any other medical condition.
- Are pregnant or plan to become pregnant.
- Are breastfeeding (nursing) or plan to breastfeed. Do not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that should not be taken with GENVOYA.

HOW TO TAKE GENVOYA

- Take GENVOYA 1 time each day with food.